

Table 10.2

Kaposi Sarcoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2011

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2009	2010-2011
SEER 9 ^b	13	26	253	805	1,479	1,994	2,102	1,922	1,674	756	319	348	262	310	269	252	246
SEER 9 except SF-Oakland	11	5	35	207	379	695	772	910	809	411	198	227	184	202	196	185	174
San Francisco-Oakland SMSA ^c	<4	21	218	598	1,100	1,299	1,330	1,012	865	345	121	121	78	108	73	67	72
San Francisco City/County	<4	18	192	519	922	1,040	1,051	748	647	237	75	75	54	73	53	49	53

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2011

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2005	2006-2007	2008-2009	2010-2011
SEER 9 ^b	0.1	0.3	2.4	7.4	13.1	16.9	17.1	14.9	12.6	5.6	2.3	2.5	1.9	2.3	2.0	1.9	1.8
SEER 9 except SF-Oakland	0.1	0.1	0.4	2.2	3.9	6.9	7.4	8.4	7.2	3.6	1.7	2.0	1.6	1.8	1.7	1.6	1.5
San Francisco-Oakland SMSA ^c	-	1.3	12.0	32.3	57.3	64.5	64.0	46.9	40.1	15.9	5.4	5.3	3.4	4.9	3.3	3.0	3.2
San Francisco City/County	-	5.3	47.6	125.1	213.9	239.5	240.8	165.5	142.3	51.8	15.9	15.9	11.4	16.1	11.5	10.5	11.3

^a Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.